

# RESOLVING THE TEMPORAL EVOLUTION OF LINE BROADENING IN SINGLE QUANTUM EMITTERS

Faculty of Mechanical Engineering, ROOM A2/201, Technická 2

**30/01/2020** 14:30

**Christian Schimpf**

*Institute of Semiconductor and Solid State Physics,  
Johannes Kepler University Linz*

Light emission from solid-state quantum emitters is inherently prone to environmental decoherence, which results in a line broadening and in the deterioration of photon indistinguishability [1]. The temporal dynamics of the dephasing mechanisms usually elude established spectroscopy methods, such as  $\mu$ PL and Michelson interferometry, due to relatively long integration times required. Photon correlation Fourier spectroscopy (PCFS) [2] allows to study the temporal evolution of line broadening in various photonic structures. The time scales we probe range from a few nanoseconds to milliseconds and, simultaneously, the spectral resolution we achieve can be as small as  $\sim 2\mu\text{eV}$ . We discuss the practical implementation of PCFS and the possibility to use it to estimate the indistinguishability of consecutively emitted single photons for applications in quantum communication and photonic-based quantum information processing [3].

[1] A. V. Kuhlmann, et al., Nat. Phys. 9, 570–575 (2013)

[2] X. Brokmann, et al., Opt. Express 14, 6333–6341 (2006)

[3] D. Huber, et al., Nat. Commun. 8, 15506 (2016)



The SINNCE project has received funding under the Horizon 2020 Research and Innovation Programme (GA 810626)

SINNCE

